

# AHCSDT407 Perform an endophytic seed test

Release: 1

## AHCSDT407 Perform an endophytic seed test

### **Modification History**

Release	TP Version	Comment
1	AHCv1.0	Initial release

# **Application**

This unit of competency describes the skills and knowledge required to determine of the presence of fungal endophyte in certain grasses.

It applies to senior seed analysts who analyse information and exercise judgement to complete a range of advanced skilled activities and demonstrate deep knowledge in a specific technical area. They have accountability for the work of others and analyse, design and communicate solutions to a range of complex problems. All work is carried out to comply with workplace procedures.

No occupational licensing, legislative or certification requirements are known to apply to this unit at the time of publication.

## Pre-requisite Unit

Nil.

#### **Unit Sector**

Seed testing (SDT)

#### **Elements and Performance Criteria**

Element	Performance criteria
Elements describe the essential outcomes.	Performance criteria describe the performance needed to demonstrate achievement of the element.
Prepare for endophyte testing	1.1 Review test request to identify samples to be tested, test methods and equipment 1.2 Clean and prepare equipment and working space 1.3 Prepare endophyte staining solution and alkaline or acid solution depending on method to be used
	1.4 Prepare working sample from Pure Seed
2. Test seed for	2.1 Soften seed for recommended time in the prepared solution at the

Approved Page 2 of 4

Element	Performance criteria
endophyte presence	prescribed temperature
	2.2 Rinse seed thoroughly in running tap water
	2.3 Remove lemma, palea and endosperm from seed and place the remaining tissue on a microscope slide in a drop of endophyte staining solution crushing if necessary
	2.4 Place cover glass on seed, applying gentle pressure and blot up excess stain
	2.5 Examine seed with compound microscope and score as positive if endophytic hyphae are present, or negative if absent
3. Test seedlings for	3.1 Take 200-250 seeds from working sample at random
endophyte presence	3.2 Sow seeds in good quality commercial potting mix at low density to allow for good growth
	3.3 Examine well-developed seedlings from the germinated sample after growing for a minimum of 42 days
	3.4 Cut seedling away from remnant seed and remove outermost sheath from the base of the seedling
	3.5 Discard sheaths which are discoloured or have developed chlorophyll
	3.6 Isolate a 3-5 mm wide longitudinal section of the sheath and place section on a microscope slide with the epidermis side down
	3.7 Stain section immediately with the endophyte staining solution for prescribed time period
	3.8 Place cover glass over the stained sheath section, press gently and blot of any excess staining solution
	3.9 Examine the section with compound microscope and score as positive or negative if endophytic hyphae are present or absent
4. Record and report test	4.1 Enter results under 'other determinations'
results	4.2 Record sample size used for testing and whether seeds or seedlings were tested
	4.3 Report results in terms of the percentage of seeds or seedlings in which endophyte was detected

# **Foundation Skills**

Foundation Skills essential to performance are explicit in the performance criteria of this unit of competency.

Approved Page 3 of 4

## **Range of Conditions**

## **Unit Mapping Information**

This unit is equivalent to AHCSDT407A Perform an endophytic seed test.

#### Links

Companion Volume implementation guides are found in VETNet - <a href="https://vetnet.gov.au/Pages/TrainingDocs.aspx?q=c6399549-9c62-4a5e-bfla-524b2322cf72">https://vetnet.gov.au/Pages/TrainingDocs.aspx?q=c6399549-9c62-4a5e-bfla-524b2322cf72</a>

Approved Page 4 of 4